

T. QIAN, Singular integrals with holomorphic kernels and Fourier multipliers on star-shaped closed Lipschitz curves ..... 193-210

G. ŁYSIK, A. Pnragmør-Littdeløw type quasi-analyticity principle ..... 217-234

E. ABON, F. GALINDO and M. LINDSTRÖM, Compact homomorphisms between algebras of analytic functions ..... 235-247

E. LELLÉVRE, Espaces BMO, inegalités de Fahey et multiplicateurs idempotents ..... 249-274

P. DŌMANSKI, J. KRONE and D. VOGR, Standard exact projective resolutions relative to a countable class of Fréchet spaces ..... 275-290

V. MÜLLER, The splitting spectrum differs from the Taylor spectrum ..... 291-294

Singular integrals with holomorphic kernels and Fourier multipliers on star-shaped closed Lipschitz curves

by

TAO QIAN (Armidale, N.S.W.)

Dedicated to Professor Alan McIntosh

STUDIA MATHEMATICA

Executive Editors: Z. Ciesielski, A. Pełczyński, W. Żelazko

The journal publishes original papers in English, French, German and Russian, mainly in functional analysis, abstract methods of mathematical analysis and probability theory. Usually 3 issues constitute a volume.

Detailed information for authors is given on the inside back cover. Manuscripts and correspondence concerning editorial work should be addressed to

STUDIA MATHEMATICA

Śniadeckich 8, P.O. Box 137, 00-950 Warszawa, Poland, fax 48-22-6293997

Abstract. The paper presents a theory of Fourier transforms of bounded holomorphic functions defined in sectors. The theory is then used to study singular integral operators on star-shaped Lipschitz curves, which extends the use of Coifman-Morrey-Rubinfeld's theory on the  $L^2$ -boundedness of the Cauchy integral operator on Lipschitz curves. The operator theory has a counterpart in Fourier multiplier theory, as well as a counterpart in function calculus of the differential operator  $\frac{d}{dz}$  on the curves.

1. Introduction. Let  $\gamma$  be a Lipschitz graph with the parameterization

$$\alpha = \alpha(t) = x + ia(t), \quad -\infty < t < \infty$$

where  $a$  is a bounded Lipschitz function,  $m = \min_{t \in \mathbb{R}} a(t)$ ,  $M = \max_{t \in \mathbb{R}} a(t)$ . Correspondence concerning subscription exchange and back numbers should be addressed to

Institute of Mathematics, Polish Academy of Sciences  
Publications Department

Śniadeckich 8, P.O. Box 137, 00-950 Warszawa, Poland, fax 48-22-6293997

$A(z) = \int_{\gamma} \frac{f(\zeta) d\zeta}{z - \zeta}$  is holomorphic in  $\{z \in \mathbb{C} : \text{Im}(z) < M - \epsilon\}$  and  $f(z) \in L^{\infty}$  on the double sectors that contain the different  $\gamma, \eta \in \gamma$  give rise to  $L^2$ -bounded Fourier multiplier operators. The Lipschitz graphs they consider are restricted to starts because of the use of a result of A. P. Calderón. The restriction can be eliminated owing to the later result of Coifman.

© Copyright by Instytut Matematyczny PAN, Warszawa, 1997

Published by the Institute of Mathematics, Polish Academy of Sciences

Printed and bound in Poland

PRINTED IN POLAND

315, 42B20; 47A60; Secondary

ISSN 0039-3223

1991 Mathematics Subject Classification: Primary 42A16



In [McQ1] the authors deal with an analogous theory on infinite Lipschitz graphs. They prove that the singular integral kernels associated with the above-mentioned  $H^\infty$ -functions are those which are holomorphic, of the Calderón-Zygmund type, and satisfy a kind of weak-boundedness condition (see (ii) of Theorem A below). In [McQ2] the converse result is proved. Another version of the theory in [McQ1], [McQ2] is the  $H^\infty$ -functional calculus of the differential operator  $\frac{1}{i} \frac{d}{dz}$ , which has also been considered for instance in [DJS] and [Mc].

Consider the holomorphic kernels on  $\mathbb{C}$ , and they should be defined on open connected sets containing  $\mathbb{R}$ . However, such holomorphic kernels with the Cauchy periodicity satisfy the standard Calderón-Zygmund size condition and are of the form  $A \cot(z/2) + \psi(z)$ , where  $A$  is a constant and  $\psi$  is a bounded holomorphic function on a bounded sector of  $\mathbb{C}$ .

The Fourier transform of  $\cot(z/2)$  is a constant multiple of the signum function (see Example (i) of §4) and the corresponding singular integral theory can be deduced, for example, from Coifman-McIntosh-Meyer's theorem [CMM].

using a partition of unity of Guy David's theory ([D]).

The second reason is related to the potential solutions of the Dirichlet and the Neumann boundary value problems on Lipschitz domains (see [FJR] and [V], for example). The star-shaped Lipschitz domains are general enough to serve this purpose, owing to the fact that every simply connected Lipschitz domain of the complex plane is the image of a star-shaped Lipschitz domain under a conformal mapping, and the fact that conformal mappings preserve harmonic functions.

The subjects presented in this paper have been further developed in various higher-dimensional cases, namely Lipschitz perturbations of the  $n$ -torus, the unit spheres of quaternionic and Clifford spaces and of the Euclidean space. They are not trivial generalizations. For instance, there

is no Liouville theorem for the  $n$ -dimensional spheres [L].

References are [Q1-5].

Special thanks are due to Ala-McIntosh and Carlos L. Kenig for their

suggestions on this topic. The author also wishes to thank Garth Gaudry,

Genfa Zhang, Joachim Hempel and Zhongmei Guo for their comments,

suggestions and their interest in this topic during my stay at the FIM,

University of Zürich, Switzerland, MITag, Leffler Institute and the New England

University, respectively. This work was partly supported by the Australian

Research Council in the form of Research Fellowship for the period when

this study was being carried out.

Finally, I refer to the Lipschitz curve defined on the interval

$[-\pi, \pi]$  with the parameterization  $f(t) = (1 + i \cot(t/2))^{-1}$ , where  $\mathbb{R}$  denotes the real number field  $\mathbb{R}$ .

The reason why we restrict ourselves to the star-shaped Lipschitz curves is as follows. Firstly, if a closed curve  $\tilde{\Gamma}$  is not star-shaped, then the corresponding difference set  $D = \{0 \neq z - \eta : z, \eta \in \tilde{\Gamma}\}$ , where

$$\tilde{\Gamma} = \left\{ \zeta = \frac{1}{i} \ln z : \operatorname{Re}(\zeta) \in [-\pi, \pi], z \in \tilde{\Gamma} \right\},$$

is not contained in any double sector defined in §2, and it may eventually spread over a region  $0 < |z| < a$ , in case  $\tilde{\Gamma}$  is winding enough. We are

considering holomorphic kernels on  $\tilde{\Gamma}$  and their generalizations in [DMS] and [McQ], the authors develop a high-dimensional theory using Clifford algebras and several complex variables which, in view of the non-commutativity of Clifford algebras, is by no means a parallel generalization of the one-dimensional case.

It is now natural to ask: is there an analogous theory for closed curves? In this paper we shall answer this question for the star-shaped Lipschitz curves given by the parameterization  $\tilde{\Gamma} = \{\exp(i\theta) : \theta \in I\}$ , where  $I = [2\pi A, 2\pi B]$ ,  $A, B \in \mathbb{R}$ ,  $A < B$ ,  $A, B \in (\pi, 3\pi]$ . It may be shown that star-shaped Lipschitz curves defined using this parameterization are the same

those defined as star-shaped and Lipschitz in the ordinary sense (see [Q3]).

One can define, in the same pattern as in the standard case, the Fourier series of  $L^2$ -functions on  $\tilde{\Gamma}$ , and the question can now be specified in the following two. The first, what kind of holomorphic kernels give rise to  $L^2$ -bounded operators on the curves? The second, is there a corresponding Fourier multiplier theory? In other words, what complex number sequences

act on  $L^2$ -functions on  $\tilde{\Gamma}$  to produce holomorphic functions on those curves? The question is non-trivial even for the case  $\tilde{\Gamma} = \mathbb{C}$ , as the Plancherel theorem does not hold in this case. On the other hand, the case  $\tilde{\Gamma} = \mathbb{R}$  is standard Calderón-Zygmund techniques (see [Z]).

The basic method of this paper is the Poisson summation formula in a sense closely related to the formula (4.1) obtained in [McQ1]. Section 2 recalls the results together with an account of some parts of the theory of Fourier transform between

the singular integral theory and the theory of the operator  $\frac{1}{i} \frac{d}{dz}$ . We also include some results on general  $L^p$ -boundedness which is an interpretation of the theory

contains some results on general  $L^p$ -boundedness. In the whole paper we shall work in the bounded case, and devote more space to those which reveal new features of the theory.



and  $\|A'\|_\infty = N < \infty$ . Denote by  $p\Gamma$  the  $2\pi$ -periodic extension of  $\Gamma$  to  $-\infty < x < \infty$ , and by  $\tilde{\Gamma}$  the closed curve

$$\tilde{\Gamma} = \{\exp(iz) : z \in \Gamma\} = \{\exp(i(x + iA(x))) : -\pi \leq x \leq \pi\}.$$

We will call  $\tilde{\Gamma}$  the *star-shaped Lipschitz curve associated with  $\Gamma$* .

We will use  $f, F$  and  $\tilde{F}$ , etc., to denote functions defined on  $p\Gamma, \Gamma$  and  $\tilde{\Gamma}$ , respectively. For  $\tilde{F} \in L^2(\tilde{\Gamma})$ , define

$$\hat{\tilde{F}}_{\tilde{\Gamma}}(n) = \frac{1}{2\pi i} \int_{\tilde{\Gamma}} z^{-n} \tilde{F}(z) \frac{dz}{z},$$

the  $n$ th Fourier coefficient of  $\tilde{F}$  with respect to  $\tilde{\Gamma}$ . We will sometimes suppress the subscript and write  $\hat{\tilde{F}}(n)$  if no confusion can occur.

Set

$$\sigma = \exp(-\max A(x)), \quad \tau = \exp(-\min A(x)).$$

Similarly to [CM1] we consider the following dense subclass of  $L^2(\tilde{\Gamma})$  (see also [GQW]):

$$A(\tilde{\Gamma}) = \{\tilde{F}(z) : \tilde{F}(z) \text{ is holomorphic in } \sigma - \eta < |z| < \tau + \eta$$

for some  $\eta > 0\}$ .

and so  $b^\pm \in H^\infty(\mathcal{S}_{\omega, \pm}^0)$ , respectively

In each of the following statements “+” should be read as either a “+”

The following transforms are used in [McQ1]

$$G^\pm(z) = c^\pm |z|^{-\frac{1}{2}} \exp(iz) b(\zeta) d, \quad z \in \mathcal{O}_{\omega, \pm}^0$$

where  $c^\pm$  is the ray length  $\mu$ ,  $\omega \in (0, \pi)$  and  $\theta$  is chosen depending on

of Cauchy's  $z \in \mathcal{O}_{\omega, \pm}^0$  so that  $\theta \in \mathcal{O}_{\omega, \pm}^0$  and  $\exp(iz)$  is exponentially decaying as

$$|z| \rightarrow \infty \text{ and}$$

complex plane

where the integral is along any path  $\delta^\pm(z)$  from  $\pm z$  to  $z$  in  $\mathcal{O}_{\omega, \pm}^0$ .

In what follows,  $c_0, c_1$ , and  $C$  will denote universal constants and  $\mathcal{O}_\mu$  will denote constants that depend on  $\omega, \mu$ , and so on, and they may vary from one occurrence to another.

Our theory is based on the main results in [McQ1] which we now reformulate for the reader's convenience.

and the sets

$$\mathcal{O}_{\omega, +}^0 = \mathcal{S}_\omega^0 \cup \{z \in \mathbb{C} : \text{Im}(z) > 0\},$$

$$\mathcal{O}_{\omega, -}^0 = \mathcal{S}_\omega^0 \cup \{z \in \mathbb{C} : \text{Im}(z) < 0\}.$$

Let  $\mathbf{X}$  be a set defined above. Denote by

$$\mathbf{X}(\pi) = \mathbf{X} \cap \{z \in \mathbb{C} : |\text{Re}(z)| \leq \pi\}$$

the truncated set, and by

$$p\mathbf{X}(\pi) = \bigcup_{k=-\infty}^{\infty} \{\mathbf{X}(\pi) + 2k\pi\}$$

the periodic set associated with the truncated one. We shall use sets of the form  $\exp(i\mathbf{O}) = \{\exp(iz) : z \in \mathbf{O}\}$ , where  $\mathbf{O}$  will be the truncated sets defined above. In the sequel  $H^\infty(\mathbf{Q})$  denotes the function space  $\{f : \mathbf{Q} \rightarrow \mathbb{C} : f \text{ is holomorphic and bounded in } \mathbf{Q}\}$ , where  $\mathbf{Q}$  will be a double or half sector defined above. We will use  $\|\cdot\|_\infty$  to denote  $\|\cdot\|_{H^\infty(\mathbf{Q})}$  if no confusion can occur.

Let  $b \in H^\infty(\mathcal{S}_\omega^0)$ ,  $\omega \in (0, \pi/2]$ . Then  $b$  can be decomposed into two parts:  $b = b^+ + b^-$ , where

$$b^+ = b\chi_{\{z: \text{Re}(z) > 0\}}, \quad b^- = b\chi_{\{z: \text{Re}(z) < 0\}},$$

Without loss of generality, we assume that  $\min A(x) <$

In this case the contours of the  $\tilde{\Gamma}$  always contain the

circle  $\mathbb{T}$ , and owing to Cauchy's theorem we have  $\hat{\tilde{F}}_{\tilde{\Gamma}}(n) = \hat{\tilde{F}}_{\mathbb{T}}(n)$ . If  $F$  &

$G$  belong to  $A(\tilde{\Gamma})$ , this remark together with Laurent-series theory imply

the Fourier inversion formula

$$\hat{\tilde{F}}_{\tilde{\Gamma}}(n) = \frac{1}{2\pi i} \int_{\tilde{\Gamma}} \tilde{F}(z) z^{-n-1} dz$$

where  $\tilde{\Gamma}$  is in the annulus in which  $\tilde{\Gamma}$  is defined, and a use of the theorem gives the Parseval formula

$$\int_{\tilde{\Gamma}} \tilde{F}(z) \overline{\tilde{G}(z)} \frac{dz}{z} = \sum_{n=-\infty}^{\infty} \hat{\tilde{F}}_{\tilde{\Gamma}}(n) \overline{\hat{\tilde{G}}_{\tilde{\Gamma}}(n)}$$

We shall use the following half and double sectors in the

$$\mathcal{O}_{\omega, \pm}^0 \text{ for } \omega \in (0, \pi/2]$$

$$\mathcal{S}_{\omega, \pm}^0 = \{z \in \mathbb{C} : |\arg(z)| < \omega, z \neq 0\}$$

$$\mathcal{S}_\omega^0 = \mathcal{S}_{\omega, +}^0 \cup \mathcal{S}_{\omega, -}^0$$





locally uniformly converges, as  $l \rightarrow \infty$ , to a  $2\pi$ -periodic and holomorphic function satisfying the assertion (i). In the sequel we shall call such sequences *applicable sequences*. Moreover, we shall show that limit functions defined through different applicable sequences differ from one another by constants bounded by  $c\|b\|_\infty$ .

To proceed, we use the decomposition

$$\begin{aligned} \sum_{k=-n}^n \phi(z + 2k\pi) &= \phi(z) + \sum_{k \neq 0}^{\pm n} (\phi(z + 2k\pi) - \phi(2k\pi)) + \sum_{k=1}^n \phi'_1(2k\pi) \\ &= \phi(z) + \sum_1 + \sum_2. \end{aligned}$$

We shall show that the series  $\sum_1$  locally uniformly converges to a bounded holomorphic function in  $S_\mu^0(\pi)$ , and some subsequence of the partial sums of  $\sum_2$  converges to a constant dominated by  $C_\mu\|b\|_\infty$ .

The convergence of  $\sum_1$  follows from the estimate

$$|\phi'(z)| \leq \frac{C_\mu}{|z|^2}, \quad z \in S_\mu^0,$$

deduced from the estimate in Corollary 1(i), the fact that  $\phi$  is holomorphic in the sectors and Cauchy's theorem. To deal with  $\sum_2$  we use the mean value theorem for integrals and we have

$$\sum_{k=1}^n \phi'_1(2k\pi) = \int_{2\pi}^{2(n+1)\pi} \phi'_1(r) dr + \sum_{k=1}^n (\phi'_1(2k\pi) - \operatorname{Re}(\phi'_1(\xi_k)) - i \operatorname{Im}(\phi'_1(\eta_k)))$$

$$\int_{-\pi}^{\pi} \phi^{\pm, \alpha}(x + 2k\pi) dx$$

$$\int_{-\pi}^{\pi} \phi^{\pm, \alpha}(\xi) dx$$

In particular, the coefficients of  $\phi^{\pm, \alpha}$  are the standard Parseval

where  $\xi_k, \eta_k \in (2k\pi, 2(k+1)\pi)$ . Owing to the estimate of  $\phi'$  again, the series in the above formula converges absolutely. The boundedness of  $\phi_1$  then guarantees the existence of an applicable sequence  $(n_l)$  such that  $\sum_{k=1}^{n_l} \phi'_1(2k\pi)$  converges to a constant  $c_0$  with the desired bound. We therefore have

$$\begin{aligned} \frac{1}{2\pi} \phi(z) &= \phi(z) + \sum_{k \neq 0} (\phi(z + 2k\pi) - \phi(2k\pi)) + \lim_{l \rightarrow \infty} \sum_{k=1}^{n_l} \phi'_1(2k\pi) \\ &= \phi(z) + \phi_0(z) + c_0, \end{aligned}$$

where  $\phi_0$  is a bounded holomorphic function in  $S_\mu^0(\pi)$  and  $c_0$  is a constant depending on the subsequence  $(n_l)$  chosen. The argument also shows that  $\phi$  can be holomorphically extended to  $\mathbf{p}S_\mu^0(\pi)$  and the different  $\phi$ 's associated with different applicable sequences may differ from one another by constants dominated by  $c\|b\|_\infty$ .

Now we prove (ii) and (iii). We use the decomposition  $b = b^+ + b^-$  indicated in §2. Define  $b^{\pm, \alpha}(z) = \exp(\mp \alpha z) b^\pm(z)$ ,  $\alpha > 0$ . Let  $\phi^\pm$  and  $\phi^{\pm, \alpha}$  be associated, according to Theorem A, with  $b^\pm$  and  $b^{\pm, \alpha}$ , respectively. Owing to the remark made after Theorem B,  $\phi^{\pm, \alpha}(\cdot) = \phi^\pm(\cdot \pm i\alpha)$ , and the latter are the inverse Fourier transforms of  $b^{\pm, \alpha}$ . We now define the corresponding holomorphic and periodic functions  $\Phi^\pm$  and  $\Phi^{\pm, \alpha}$  in  $\mathbf{p}C_{\omega, \pm}^0(\pi)$ , respectively, which satisfy the size condition in the assertion (i). It is to be noted that for all  $\Phi^{\pm, \alpha}$  we may, and we actually do, choose the same applicable sequence  $(n_l)$  for  $\Phi^{\pm, \alpha}$  as we have chosen for  $\Phi^\pm$ . Using the estimate in Corollary 1(i) and the fact that  $\phi$  is holomorphic, we can show that the convergence of  $\sum_1$  is locally (in  $z$ ) uniform for  $\alpha \rightarrow 0$ , and is absolute. Let

$$\begin{aligned} \frac{1}{2\pi} \Phi^{\pm, \alpha}(z) &= \phi^{\pm, \alpha}(z) + \phi_0^{\pm, \alpha}(z) + c_0^{\pm, \alpha}, \\ \frac{1}{2\pi} \Phi^\pm(z) &= \phi^\pm(z) + \phi_0^\pm(z) + c_0^\pm, \end{aligned}$$

where  $\phi_0^{\pm, \alpha}$  and  $\phi_0^\pm$  are holomorphic and uniformly (for  $\alpha \rightarrow 0$ ) bounded in  $C_{\mu, \pm}^0(\pi)$ . Since the convergence as  $n_l \rightarrow \infty$  is uniform for  $\alpha \rightarrow 0$ , we can exchange the order of taking the limits as  $n_l \rightarrow \infty$  and  $\alpha \rightarrow 0$ , and conclude that  $\phi^{\pm, \alpha}$ ,  $\phi_0^{\pm, \alpha}$  and  $c_0^{\pm, \alpha}$  converge to  $\phi^\pm$ ,  $\phi_0^\pm$  and  $c_0^\pm$ , respectively, locally uniformly in  $C_{\omega, \pm}^0(\pi)$ . Therefore,  $\lim_{\alpha \rightarrow 0} \Phi^{\pm, \alpha}(z) = \Phi^\pm(z)$ . Since for a fixed  $\alpha$ ,  $\Phi^{\pm, \alpha} \in L^\infty([-\pi, \pi])$ , and the series which defines  $\Phi^{\pm, \alpha}$  converges uniformly in  $x \in [-\pi, \pi]$  as  $n_l \rightarrow \infty$ , we have

$$\int_{-\pi}^{\pi} \exp(-i\xi x) \Phi^{\pm, \alpha}(x) dx = \int_{-\pi}^{\pi} \exp(-i\xi x) \lim_{l \rightarrow \infty} \sum_{k=1}^{n_l} \phi^{\pm, \alpha}(x + 2k\pi) dx = \int_{-\pi}^{\pi} \exp(-i\xi x) \phi^{\pm, \alpha}(x) dx$$

for all non-zero real  $\xi$  in the sense of (3) and  $\{b^{\pm, \alpha}(n)\}$ ,  $n \neq 0$ , are the standard Fourier coefficients of any smooth and periodic function on  $[-\pi, \pi]$ , the identity holds.

$$\int_{-\pi}^{\pi} \exp(-i\xi x) \phi^{\pm, \alpha}(x) dx = \sum_{n=-\infty}^{\infty} b^{\pm, \alpha}(n) \hat{F}^{\pm, \alpha}(\xi)$$

Proceed as in [Mc],

on letting  $\alpha \rightarrow 0$  we

$\phi$  can be holomorphically extended to  $\mathbf{p}S_\mu^0(\pi)$  and the different  $\phi$ 's associated with different applicable sequences may differ from one another by constants dominated by  $c\|b\|_\infty$ .

where  $\hat{F}^{\pm, \alpha}(\xi) = \int_{-\pi}^{\pi} \exp(-i\xi x) \phi^{\pm, \alpha}(x) dx$  and  $\hat{F}^\pm(\xi) = \int_{-\pi}^{\pi} \exp(-i\xi x) \phi^\pm(x) dx$ .

Let  $\xi > 0$ . Since  $F(n)$  decays rapidly as  $n \rightarrow \infty$





integral over the last mentioned contour is bounded, using only the fact that  $\pm \text{Re}(z) > 0$ . Therefore  $b$  is well defined with the desired bounds. We leave the details to the interested reader (or refer to [Q4]).

Let  $F$  be any  $2\pi$ -periodic function on  $\mathbb{T}$ . Expanding  $F$  in Fourier series and using the definition of  $\hat{b}_\varepsilon$ , we have

$$2\pi \sum_{n=-\infty}^{\infty} \hat{b}_\varepsilon(n) \hat{F}_{[-\pi, \pi]}(-n) = \int_{\varepsilon < |x| < \pi} \Phi(x) F(x) dx - \Phi_1(\varepsilon) F(0).$$

On letting  $\varepsilon \rightarrow 0$ , we get

$$2\pi \sum_{n=-\infty}^{\infty} b(n) \hat{F}_{[-\pi, \pi]}(-n) = \lim_{\varepsilon \rightarrow 0} \left\{ \int_{\varepsilon < |x| < \pi} \Phi(x) F(x) dx - \Phi_1(\varepsilon) F(0) \right\}.$$

Denoting by  $(G(b), G_1(b))$  a pair of holomorphic functions associated with  $b$  in the pattern of Theorem 1, from the Parseval identity it follows that

$$\lim_{\varepsilon \rightarrow 0} \left\{ \int_{\varepsilon < |x| < \pi} (G(b)(x) - \Phi(x)) F(x) dx + (G_1(\varepsilon) - \Phi_1(\varepsilon)) F(0) \right\} = 2\pi(b_1(0) - b(0)) \hat{F}_{[-\pi, \pi]}(0),$$

where  $b_1(0)$  is associated with  $(G(b), G_1(b))$  in the Parseval identity (iii) of Theorem 1. According to Theorem 1 (see also the argument at the end of its proof), we can add any constant to  $G(b)$  and accordingly adjust the value of  $b_1(0)$  in order to make (iii) of Theorem 3 still hold. In particular, we can choose a constant such that  $b_1(0) - b(0) = 0$ . The right hand side of the last displayed equality then becomes zero. Using an approximation to identity  $(F_n)$  with the property  $F_n(0) = 0$  for all  $n$ , we conclude that  $G(b)(x) = \Phi(x)$  for  $x \neq 0$ , which implies  $G(b)(z) = \Phi(z)$  for all  $z \in \mathbb{S}_\omega^0(\pi)$  owing to analyticity. Using the assertion (ii) of Theorem 1 on  $G_1(b)$  and the assumption (iii) on the function  $\Phi$ , we have  $G_1(b) = \Phi_1$  and  $b^\pm = 0$ .

According to Theorem 3 is identical. The uniqueness of  $b$  can be proved similarly. The proof is complete.

**4. Singular integrals on star-shaped Lipschitz curves.** The results obtained in §3 can be used to study the relations between singular integrals and multiplier transforms on periodic Lipschitz curves. Alternatively we can consider the closed star-shaped Lipschitz curves defined in §2. By performing the change of variable  $z \rightarrow \exp(iz)$  and substituting  $\tilde{\Phi} = \Phi \circ (\frac{1}{i} \ln)$  and  $\tilde{\Phi}_1 = \Phi_1 \circ (\frac{1}{i} \ln)$  in Theorems 1 and 2, we obtain the following the-

**THEOREM 3.** Let  $\omega \in (0, \pi/2]$  and  $b \in H^\infty(\mathbb{S}_\omega^0)$ . Then there exists a pair of functions  $(\tilde{\Phi}, \tilde{\Phi}_1)$  such that  $\tilde{\Phi}$  and  $\tilde{\Phi}_1$  are holomorphic in  $\exp(i\mathbb{S}_\omega^0(\pi))$  and  $\exp(i\mathbb{S}_{\omega,+}^0(\pi))$ , respectively and for every  $\mu \in (0, \omega)$ ,

$$(i) \quad \|\tilde{\Phi}(z)\| \leq \frac{C_{\omega, \mu} \|b\|_\infty}{1 - |z|}, \quad z \in \exp(i\mathbb{S}_\omega^0(\pi)),$$

$$(ii) \quad \tilde{\Phi}_1 \in H^\infty(\exp(i\mathbb{S}_\mu^0(\pi))), \quad \|\tilde{\Phi}_1\|_{H^\infty(\exp(i\mathbb{S}_\mu^0(\pi)))} \leq C_{\omega, \mu} \|b\|_\infty,$$

$$\tilde{\Phi}'_1(z) = \frac{1}{iz} (\tilde{\Phi}(z) + \tilde{\Phi}(z^{-1})), \quad z \in \exp(i\mathbb{S}_{\omega,+}^0(\pi))$$

$$(iii) \quad 2\pi \sum_{n=-\infty}^{\infty} b(n) \hat{F}_\mathbb{T}(-n) = \lim_{\varepsilon \rightarrow 0} \left\{ \int_{|\ln z| > \varepsilon, z \in \mathbb{T}} \tilde{\Phi}(z) \tilde{F}(z) \frac{dz}{z} + \tilde{\Phi}_1(\exp(i\varepsilon)) \tilde{F}(0) \right\}$$

for all smooth functions  $\tilde{F}$  defined on  $\mathbb{T}$ , where  $\hat{F}_\mathbb{T}(n)$  is the  $n$ th Fourier coefficient of  $\tilde{F}$  and  $b(0) = \frac{1}{2\pi} \tilde{\Phi}_1(\exp(i\pi))$ .

**THEOREM 4.** Let  $\omega \in (0, \pi/2]$  and  $(\tilde{\Phi}, \tilde{\Phi}_1)$  be a pair of holomorphic functions defined in  $\exp(i\mathbb{S}_\omega^0(\pi))$  and  $\exp(i\mathbb{S}_{\omega,+}^0(\pi))$ , respectively, satisfying

(i) there is a constant  $c_0$  such that

$$\|\tilde{\Phi}(z)\| \leq \frac{c_0}{|1 - z|}, \quad z \in \exp(i\mathbb{S}_\omega^0(\pi));$$

(ii) there is a constant  $c_1$  such that  $\|\tilde{\Phi}_1\|_{H^\infty(\exp(i\mathbb{S}_{\omega,+}^0(\pi)))} < c_1$ , and

$$\tilde{\Phi}'_1(z) = \frac{1}{iz} (\tilde{\Phi}(z) + \tilde{\Phi}(z^{-1})), \quad z \in \exp(i\mathbb{S}_{\omega,+}^0(\pi)).$$

Then for every  $\mu \in (0, \omega)$ , there exists a function  $b^\mu$  in  $H^\infty(\mathbb{S}_\mu^0)$ ,

and the function pair determined by  $b^\mu$  according to  $(\tilde{\Phi}, \tilde{\Phi}_1)$  modulo additive constants. Moreover,

$$b^\pm(n) = \frac{1}{2\pi} \lim_{\varepsilon \rightarrow 0} \int_{-i \ln z \in A^\pm(\varepsilon, \theta, \varrho)} z^{-n} \tilde{\Phi}(z) dz$$

where the contour  $A^\pm(\varepsilon, \theta, \varrho)$  is defined in

$$\tilde{\Phi}_1(\exp(iz)) = \frac{1}{i} \tilde{\Phi}'_1(z)$$

where  $(z)$  is any path from



The following corollaries are versions of Theorems 3 and 4 in terms of holomorphic extension of series of positive and negative powers (see also the paragraph following the statement of Theorem 6 below).

COROLLARY 2. Let  $(b_n)_{n=\pm 1}^{\pm\infty} \in l^\infty$  and  $\tilde{\Phi}(z) = \sum_{n=\pm 1}^{\pm\infty} b_n z^n$ ,  $|z^{\pm 1}| < 1$ , and  $\omega \in (0, \pi/2)$ . If there exist  $\delta > 0$  such that  $\omega + \delta \leq \pi/2$ , and a function  $b \in H^\infty(S_{\omega+\delta, \pm}^0)$  such that  $b(n) = b_n$  for all  $\pm n = \pm 1, \pm 2, \dots$ , then the function  $\tilde{\Phi}$  can be holomorphically extended to the region  $\exp(iC_{\omega+\delta, \pm}^0)$ .

Moreover, we have

$$|\tilde{\Phi}(z)| \leq \frac{C_{\omega, 0}}{|Fz|} \quad z \in \exp(iC_{\omega, \pm}^0)$$

COROLLARY 3. Let  $\omega \in (0, \pi/2)$ , and  $\tilde{\Phi}$  be holomorphic and satisfy

$$|\tilde{\Phi}(z)| \leq \frac{C_{\omega, 0}}{|Fz|} \quad z \in \exp(iC_{\omega, \pm}^0)$$

Then for every  $u \in (0, \omega)$ , there exists a function  $b^u$  such that  $b^u \in H^\infty(S_{\omega-u, \pm}^0)$

and  $\tilde{\Phi}(z) = \sum_{n=\pm 1}^{\pm\infty} b^u(n) z^n$ . Moreover,  $b^u = \lim_{\varepsilon \rightarrow 0+} b^\varepsilon$

$$b^\varepsilon(n) = \lim_{\varepsilon \rightarrow 0+} \frac{1}{2\pi} \int_{\gamma_\varepsilon} \exp(-in\varepsilon) \tilde{\Phi}(\exp(i\varepsilon z)) dz = \tilde{\Phi}(b^\varepsilon(n))$$

where  $A^\pm(\varepsilon, \theta, \rho)$  is defined as in Theorem 2, and

$$\tilde{\Phi}_1(\exp(i\varepsilon)) = \int \tilde{\Phi}(\exp(iz)) dz$$

where  $\gamma_\varepsilon$  is any path from  $-\varepsilon$  to  $\varepsilon$  lying in  $C_{\omega, \pm}^0$ .

Proof.  $\tilde{\Phi}(z) = \sum_{n=\pm 1}^{\pm\infty} b_n z^n$  is not single-valued. In fact, if  $\mu_1 \neq \mu_2$ , then both  $b^{\mu_1}$  and  $b^{\mu_2}$  satisfy the requirement, but  $b^{\mu_1} \neq b^{\mu_2}$  in general. This can be verified by using  $\tilde{\Phi}(z) = z^n$ ,  $n \in \mathbb{Z}$ , for example (see also [Q4]).

Each time, remember to observe the following application of Corollary 2.

COROLLARY 4. Let  $\omega \in (0, \pi/2)$ , where  $\omega$  is not a rational multiple of  $\pi$ , and let  $\tilde{\Phi}$  be holomorphic and satisfy

$$|\tilde{\Phi}(z)| \leq \frac{C_{\omega, 0}}{|Fz|} \quad z \in \exp(iC_{\omega, \pm}^0)$$

Then  $\tilde{\Phi}(z) = 0$  for all  $z$  in the region  $\exp(iC_{\omega, \pm}^0)$ .

Proof. It is well known that  $\tilde{\Phi}$  does not have any holomorphic extension across intervals on the unit circle, and hence, according to Corollary 2, it is not holomorphic in any region  $S_{\omega+\delta, \pm}^0$  in the pattern of Corollary 2.

For a function  $b$  and a function  $\tilde{F}$  defined in Theorem 3, by Laurent series theory, the series

$$\sum_{n=-\infty}^{\infty} b(n) \tilde{F}_T(n) z^n$$

locally uniformly converges to a holomorphic function in the annulus on which  $\tilde{F}$  is defined. Recalling the fact that  $\tilde{F}_T(n) = \tilde{F}(z) = \exp(i\theta n)$ , we define an operator  $M_b : A(\tilde{F}) \rightarrow A(\tilde{F})$  by

$$M_b F(z) = 2\pi \sum_{n=-\infty}^{\infty} b(n) \tilde{F}_T(n) z^n$$

On the other hand, for a pair of functions  $(\tilde{\Phi}, \tilde{\Phi}_1)$  specified in Theorem 3,

$$\tilde{\Phi}_1(\exp(i\varepsilon)) = \int_{\gamma_\varepsilon} \tilde{\Phi}(\exp(iz)) dz$$

where  $\gamma_\varepsilon$  is the normalized tangent vector to  $A^\pm(\varepsilon, \theta, \rho)$  at  $\varepsilon$  lying in  $C_{\omega, \pm}^0$ . We have the following theorem.

THEOREM 5. Let  $\omega \in (\arctan \lambda, \pi/2)$ ,  $b \in H^\infty(S_{\omega, \pm}^0)$  and  $(\tilde{\Phi}, \tilde{\Phi}_1)$  be the function pair associated with  $\omega$  in the pattern of Theorem 3. Then

- (i)  $T_b \tilde{\Phi}_1$  is a well-defined operator from  $A(\tilde{F})$  to  $A(\tilde{F})$ , and  $M_b$  modulo a constant multiple of the identity operator;
- (ii)  $M_b$  extends to a bounded operator on  $L^2(\Gamma)$ , whose operator norm is  $\|b\|_\infty$ .

Proof. (i) For any  $\alpha > 0$ , define  $b^{\pm\alpha}(z) = z^\alpha b(z)$ ,  $b^{\pm\alpha}(n) = z^\alpha b(n)$  are the functions defined in the proof of Theorem 1. Let  $(\tilde{\Phi}^{\pm\alpha}, \tilde{\Phi}_1^{\pm\alpha})$  be the function pair associated with  $b^{\pm\alpha}$  in the pattern of Theorem 3. To verify (i) and (ii), we have

$$M_b F(z) = 2\pi \sum_{n=-\infty}^{\infty} b(n) \tilde{F}_T(n) z^n$$

Taking the limit  $\alpha \rightarrow 0$  as in the proof of Theorem 1, we have

$$\tilde{\Phi}_1^{\pm\alpha}(z) = \tilde{\Phi}_1(z)$$

Obtain the desired equality for  $b$  and hence for  $M_b$ .





(ii) One can alternatively prove the boundedness of the operator

$$T_{(\Phi, \Phi_1)} F(z) = \lim_{\varepsilon \rightarrow 0} \left\{ \int_{\pi \geq |\operatorname{Re}(z-\eta)| > \varepsilon} \Phi(z-\eta) F(\eta) d\eta + \Phi_1(\varepsilon t(z)) F(z) \right\},$$

$$F \in \mathcal{A}(\Gamma),$$

where  $t(z)$  is the normalized tangent vector of  $\Gamma$  at  $z$  lying inside  $S_{\omega,+}^0(\pi)$ , and  $\mathcal{A}(\Gamma)$  is the class of  $2\pi$ -periodic and holomorphic functions defined by the condition  $F \in \mathcal{A}(\Gamma)$  if and only if  $\tilde{F} = F \circ (i^{-1} \ln) \in \mathcal{A}(\tilde{\Gamma})$ . Owing to the decomposition of  $\Phi$  in the assertion (i) of Theorem 1, we have

$$T_{(\Phi, \Phi_1)} F(z) = \lim_{\varepsilon_n \rightarrow 0} \left\{ \int_{\pi \geq |\operatorname{Re}(z-\eta)| > \varepsilon_n} \phi(z-\eta) F(\eta) d\eta + \int_{\pi \geq |\operatorname{Re}(z-\eta)| > \varepsilon_n} \phi_0(z-\eta) F(\eta) d\eta \right\} + c_1 \int_{-\pi}^{\pi} F(\eta) d\eta + c_2 F(z),$$

where  $\varepsilon_n \rightarrow 0$  is an appropriate subsequence of  $\varepsilon \rightarrow 0$ , and  $c_1$  and  $c_2$  are constants

The second and the third integrals are dominated by the  $L^2$ -norm of  $F$ , while the first integral is dominated by

$$\sup_{\varepsilon > 0} \left| \int_{|\operatorname{Re}(z-\eta)| > \varepsilon} \phi(z-\eta) F_1(\eta) d\eta \right| + c M F_1(z) \quad \operatorname{Re}(z) \in ]-\pi, \pi[$$

wise, and  $M F_1$  is the surface Dirac operator on every star-shaped Lipschitz curve. If  $\operatorname{Im}(\lambda) > 0$ , then from [McQ], we have

$$\phi_\lambda(z) = \begin{cases} \exp(i\lambda z) & \text{if } \operatorname{Re}(z) > 0 \\ 0 & \text{if } \operatorname{Re}(z) < 0 \end{cases}$$

If  $\operatorname{Im}(\lambda) < 0$ , then we have

$$\phi_\lambda(z) = \begin{cases} 0 & \text{if } \operatorname{Re}(z) > 0 \\ \exp(i\lambda z) & \text{if } \operatorname{Re}(z) < 0 \end{cases}$$

We state without proof the following theorem. For a proof we refer the reader to [McQ2].

**THEOREM 6** Let  $\omega \in ]\arctan N, \pi/2[$ ,  $\Phi$  be holomorphic in  $\exp(iS_{\omega,+}^0)$  and satisfy (4) of Theorem 4 with respect to  $\omega$ . If  $T$  is a bounded operator on  $L^2(\tilde{\Gamma})$  and

$$T(\tilde{F})(z) = \int_{\tilde{\Gamma}} \tilde{\Phi}(z\zeta^{-1}) \tilde{F}(\zeta) \frac{d\zeta}{\zeta}, \quad z \notin \operatorname{supp}(\tilde{F}),$$

for all  $\tilde{F} \in C_0(\tilde{\Gamma})$ , the class of continuous functions, then there exists a unique function  $\tilde{\Phi}_1 \in H^\infty(\exp(iS_{\omega,+}^0))$ ,  $\mu \in (0, \omega)$ , such that

$$\tilde{\Phi}_1'(z) = \frac{1}{iz} (\tilde{\Phi}(z) + \tilde{\Phi}(z^{-1})), \quad z \in \exp(iS_{\omega,+}^0(\pi)),$$

and

$$T(\tilde{F}) = T_{(\tilde{\Phi}, \tilde{\Phi}_1)}(\tilde{F})$$

for all  $\tilde{F} \in C_0(\tilde{\Gamma})$ .

As stated in Corollary 2, for  $b \in S_{\omega}^0$  the function  $\tilde{\Phi}^+(z) = \sum_{n=1}^{\infty} b(n)z^n$ ,  $|z| < 1$ , can be holomorphically extended to  $\exp(iC_{\omega,+}^0(\pi))$ , and  $\tilde{\Phi}^-(z) = \sum_{n=-\infty}^{-1} b(n)z^n$ ,  $|z| > 1$ , can be holomorphically extended to  $\exp(C_{\omega,-}^0(\pi))$ . So, we have the expression

$$(6) \quad \tilde{\Phi}(z) = \sum_{n=-\infty}^{\infty} b(n)z^n, \quad z \in \exp(iS_{\omega}^0(\pi)).$$

In many cases using (6) is more convenient than using (4) in finding an explicit formula for  $\tilde{\Phi}$ , and hence for  $\Phi$ .

**EXAMPLE (1)** Let  $\phi(z) = -i \operatorname{sgn}(z)$ , then from [McQ], we get  $\phi(z) = \frac{1}{z} - \frac{1}{z^{-1}}$ , which corresponds to the Hilbert transform with kernel  $\phi(z)$ . Using the expression (6), we obtain  $\tilde{\Phi}(z) = \cot \frac{z}{2}$ ,  $\tilde{\Phi}_1(z) = -i \frac{1+z}{1-z}$ , and  $\tilde{\Phi}_1'(z) = \frac{2}{z^2}$ . From the assertion (i) of Theorem 5, the Fourier multiplier  $-i \operatorname{sgn}$  corresponds to the kernels  $\frac{z^{-1}}{2\pi} \cdot \frac{1+z}{1-z}$  and  $\frac{z}{2\pi} \cot \frac{z}{2}$  on  $\tilde{\Gamma}$  and  $\Gamma$ , respectively

**EXAMPLE (2)** where  $\phi(z) = \frac{1}{z} - \frac{1}{z^{-1}}$ , we get  $\tilde{\Phi}(z) = \frac{1}{z} - \frac{1}{z^{-1}}$  and  $\tilde{\Phi}_1(z) = 0$ . The Hardy-Littlewood maximal operators of  $F_1$  on the  $c$ -boundedness results for the operator introduced by  $\phi$ .  $\phi$  for the Hardy-Littlewood maximal function, we obtain  $c$ -boundedness.

**Remark** There is some interest in direct proofs for Theorem 6, which we refer to [GQW] and [Q2].

It is easy to see that in each of the two cases  $\phi_\lambda$  is in  $L^1(\mathbb{R}) \cap L^2(\mathbb{R})$  and so the remark we made after Theorem 5 applies to both cases.

From the definition, for  $\operatorname{Im}(\lambda) > 0$ , we have

$$\Phi_\lambda(z) = \begin{cases} \frac{i \exp(i\lambda(z+2\pi))}{1 - \exp(i\lambda 2\pi)} & \text{if } -\pi < \operatorname{Re}(z) < 0 \\ \frac{i \exp(i\lambda z)}{\exp(i\lambda 2\pi)} & \text{if } 0 < \operatorname{Re}(z) < \pi \end{cases}$$



For  $\text{Im}(\lambda) < 0$ ,

$$\Phi_\lambda(z) = \begin{cases} \frac{-i \exp(i\lambda(z - 2\pi))}{1 - \exp(-i\lambda 2\pi)} & \text{if } 0 < \text{Re}(z) < \pi, \\ \frac{-i \exp(i\lambda z)}{1 - \exp(-i\lambda 2\pi)} & \text{if } -\pi < \text{Re}(z) < 0. \end{cases}$$

For  $\text{Im}(\lambda) > 0$ ,

$$\tilde{\Phi}_\lambda(z) = \begin{cases} \frac{i \exp(i\lambda 2\pi) z^\lambda}{1 - \exp(i\lambda 2\pi)} & \text{if } -\pi < \text{Re}\left(\frac{\ln z}{i}\right) < 0, \\ \frac{iz^\lambda}{1 - \exp(i\lambda 2\pi)} & \text{if } 0 < \text{Re}\left(\frac{\ln z}{i}\right) < \pi. \end{cases}$$

For  $\text{Im}(\lambda) < 0$ ,

$$\tilde{\Phi}_\lambda(z) = \begin{cases} \frac{-i \exp(-i\lambda 2\pi) z^\lambda}{1 - \exp(-i\lambda 2\pi)} & \text{if } 0 < \text{Re}\left(\frac{\ln z}{i}\right) < \pi, \\ \frac{-i \exp(-i\lambda z)}{1 - \exp(-i\lambda 2\pi)} & \text{if } -\pi < \text{Re}\left(\frac{\ln z}{i}\right) < 0. \end{cases}$$

As in the above examples,  $\frac{1}{2\pi}$  times the above functions will be the kernels of the resolvents on the curves  $\tilde{\Gamma}$  and  $\tilde{\Gamma}_1$  respectively.

We now outline how the  $H^\infty$ -functional calculus developed in [Mc] can be applied to the present case (see [McQ1], [McQ3] for the infinite Lipschitz graph case), and indicate the relation between this functional calculus and the operator classes  $\tilde{M}_h$  and  $T_{(\tilde{\Phi}, \tilde{\Phi}_1)}$ .

For a function  $\tilde{F} \in A(\tilde{\Gamma})$  we define the differential operator  $\frac{d}{dz}|_{\tilde{\Gamma}}$  by

$$\frac{d}{dz}|_{\tilde{\Gamma}} \tilde{F}(z) = \lim_{\substack{h \rightarrow 0 \\ z+h \in \tilde{\Gamma}}} \frac{\tilde{F}(z+h) - \tilde{F}(z)}{h} \quad \text{for } z \in \tilde{\Gamma}.$$

For  $1 < p < \infty$ ,  $(L^p(\tilde{\Gamma}), L^{p'}(\tilde{\Gamma}))$  is a pairing of Banach spaces given by  $\langle \tilde{F}, \tilde{G} \rangle = \int_{\tilde{\Gamma}} \tilde{F}(z) \tilde{G}(z) dz$ , where  $p' = (1 - p^{-1})^{-1}$ . Now use duality to define  $D_{\tilde{\Gamma}, p}$  as the closed operator with the largest domain in  $L^p(\tilde{\Gamma})$  which satisfies

$$\langle D_{\tilde{\Gamma}, p} \tilde{F}, \tilde{G} \rangle = \left\langle \tilde{F}, -z \frac{d}{dz} \Big|_{\tilde{\Gamma}} \tilde{G} \right\rangle$$

for all  $\tilde{F}$  and  $\tilde{G}$  in  $A(\tilde{\Gamma})$ .

Let  $\omega \in (\arctan N, \pi/2]$  and  $\lambda \notin \mathbf{S}_\omega^0$ . It is easy to verify that  $D_{\tilde{\Gamma}, p}$  is the surface Dirac operator on  $\tilde{\Gamma}$  and the function  $\frac{1}{2\pi} \tilde{\Phi}_\lambda$  given in Example (ii) is the convolution kernel of the resolvent operator  $(D_{\tilde{\Gamma}, p} - \lambda)^{-1}$  in the sense

of Theorem 5. Moreover,

$$\begin{aligned} \|(D_{\tilde{\Gamma}, p} - \lambda)^{-1}\| &\leq \left\| \frac{1}{2\pi} \tilde{\Phi}_\lambda \right\|_{L^1(\tilde{\Gamma})} \leq \sum_{n=-\infty}^{\infty} \|\phi_\lambda(\cdot + 2\pi n)\|_{L^1(\Gamma)} \\ &= \|\phi_\lambda\|_{L^1(p\Gamma)} \leq \sqrt{1 + N^2} \{\text{dist}(\lambda, \mathbf{S}_\omega^0)\}^{-1}, \end{aligned}$$

where we have used the bounds of  $\|\phi_\lambda\|_{L^1(p\Gamma)}$  obtained in [McQ1].

The above estimate implies that  $D_{\tilde{\Gamma}, p}$  is a *type- $\omega$*  operator ([Mc]) that allows us to define  $b(D_{\tilde{\Gamma}, p})$  via spectral integrals first for those  $H^\infty$ -functions  $b$  with good decay properties at both 0 and  $\infty$ :

$$b(D_{\tilde{\Gamma}, p}) = \frac{1}{2\pi i} \int_\delta b(\eta) (D_{\tilde{\Gamma}, p} - \eta I)^{-1} d\eta,$$

where  $\delta$  is a path consisting of four rays:  $\{s \exp(-i\theta) : s \text{ goes from } \infty \text{ to } 0\}$ ,  $\{s \exp(i\theta) : s \text{ goes from } 0 \text{ to } \infty\}$ ,  $\{s \exp(i(\pi + \theta)) : s \text{ goes from } \infty \text{ to } 0\}$ ,  $\{s \exp(i(\pi - \theta)) : s \text{ goes from } 0 \text{ to } \infty\}$ , where  $\arctan N < \theta \leq \omega$ .

It is not difficult to show using the above estimate that each  $b \in \mathcal{H}_{\tilde{\Gamma}, p}$  is a bounded operator, and  $b(D_{\tilde{\Gamma}, p}) = \tilde{M}_b \equiv T_{(\tilde{\Phi}, \tilde{\Phi}_1)}$ . Taking limits of sequences of Calderón-Zygmund operators (see [Mc] or [GM2]), one can then extend the definition of  $b(D_{\tilde{\Gamma}, p})$  to all the functions in  $H^\infty(\mathbf{S}_\omega^0)$ , and prove

$$b(D_{\tilde{\Gamma}, p}) = \tilde{M}_b \equiv T_{(\tilde{\Phi}, \tilde{\Phi}_1)},$$

with the properties

$$\|b(D_{\tilde{\Gamma}, p})\| \leq C_\omega \|b\|_\infty,$$

$$(b_1 b_2)(D_{\tilde{\Gamma}, p}) = b_1(D_{\tilde{\Gamma}, p}) b_2(D_{\tilde{\Gamma}, p}),$$

$$(\alpha_1 b_1 + \alpha_2 b_2)(D_{\tilde{\Gamma}, p}) = \alpha_1 b_1(D_{\tilde{\Gamma}, p}) + \alpha_2 b_2(D_{\tilde{\Gamma}, p}),$$

whenever  $b_1, b_2 \in H^\infty(\mathbf{S}_\omega^0)$  and  $\alpha_1, \alpha_2$  are complex numbers.

**5. Fourier multipliers on star-shaped Lipschitz curves.** In this section we shall not restrict ourselves to the  $H^\infty$ -multipliers. We will point out that all the results and methods of the Fourier multiplier theory for the infinite Lipschitz graph case developed in [McQ3] can be adapted to the present case. The major changes are that the class  $A(\tilde{\Gamma})$  is replaced by  $A(\Gamma)$  for our purpose, and whenever we deal with a kernel on  $\Gamma$  we refer to the corresponding kernel on  $p\Gamma$  via the Poisson summation formula. We state two results without proofs. Both can be proved using the corresponding Schur lemma in the present case (see [McQ3]).



For  $b = (b_n)_{n \geq 0} \in l^\infty$ , define

$$\|b\|_{M_p(\Gamma)} = \sup \left\{ \left\| \sum_{n=0}^{\infty} b_n F(n)z^n \right\|_{L^p(\Gamma)} : \|F\|_{L^1(\Gamma)} \leq 1 \right\}$$

$$M_p(\tilde{\Gamma}) = \{b : \|b\|_{M_p(\tilde{\Gamma})} < \infty\}$$

Elements  $b$  in  $M_p(\tilde{\Gamma})$  are called  $L^p(\tilde{\Gamma})$ -Fourier multipliers

[Aster] Asterisque 57 (1978) -  
 ever. L'intégrale de Cauchy définie sur  
 des lipschitziennes. Ann. of Math. 116  
 liers sur certaines courbes au plan com-  
 1984), 157-186.  
 185) Opérateurs de Calderón-Zygmunda  
 on. Rev. Mat. Iberoamericana - (1985).

[M. Riviere] Potential techniques for  
 ns, Acta Math. 141 (1978), 163-186.  
 [Boundedness of singular integral oper-  
 -shaped Lipschitz-curves, Colloq. Math.

afford algebras, Fourier transforms and  
 hix surfaces. Rev. Mat. Iberoamericana  
 [McM3] A. McIntosh and S. Semmes - Convolution singular integral operators  
 on Lipschitz surfaces. Illinois Math. Surv. 5 (1992), 45-48.  
 [McM4] A. McIntosh - Operators which have an  $H^\infty$ -functional calculus. in: Mimico  
 and Pagan. Dublin: Academic Press, 1986, 2-10.  
 [Q1] A. McIntosh and T. Qian - Convolution singular integral operators on Lip-  
 schitz curves. in: Lecture Notes in Math. 1494, Springer, 1991, 142-162.  
 [McQ2] Singular integrals along Lipschitz curves with holomorphic kernels.  
 Proc. Amer. Math. Soc. 119 (1993), 49-57.  
 [McQ3] Singular integrals along Lipschitz curves with holomorphic kernels.  
 Proc. Amer. Math. Soc. 121 (1994), 167-176.

[Q1] Singular integrals on the  $n$ -cube with holomorphic kernels.  
 Proc. Amer. Math. Soc. 121 (1994), 94-100.

[Q2] Transference from Lipschitz graphs to periodic Lipschitz graphs. in: Mimico  
 and Pagan. Dublin: Academic Press, 1986, 2-10.

[Q3] Singular integrals on star-shaped Lipschitz surfaces in the quaternionic  
 space, preprint.

otherwise. Using  $F(z) = \frac{1}{1 - \exp(z)}$  one can show that  
any  $z \in \Gamma$ .

[CM1] R. Coifman and A. Meyer - Fourier analysis of multilinear convolution  
 operators. Stud. Adv. Math. 19 (1982), 61-74.

satisfying  $\Phi(r \exp(i\theta)) = \psi(\exp(i\theta))$ , where  $\int_{-\pi}^{\pi} \psi(\exp(i\theta)) d\theta < \infty$ . Then  
 $b = (\hat{\Phi}(n))_{n \geq 0} \in M_p(\tilde{\Gamma})$ ,  $1 < p < \infty$  and the associated convolution  
 operator  $T_b$  is given by

$$T_b \tilde{F}(z) = \int_{\tilde{\Gamma}} \hat{\Phi}(zn^{-1}) \tilde{F}(\eta) \frac{d\eta}{\eta}, \quad \tilde{F} \in \mathcal{A}(\tilde{\Gamma}).$$

Let  $\tilde{\Gamma}_1$  and  $\tilde{\Gamma}_2$  be two curves of the type under consideration. Define

$$M_p(\tilde{\Gamma}_1, \tilde{\Gamma}_2) = \{b \in l^\infty : \|b\|_{M_p(\tilde{\Gamma}_1, \tilde{\Gamma}_2)} < \infty\},$$

where

$$\|b\|_{M_p(\tilde{\Gamma}_1, \tilde{\Gamma}_2)} = \sup \left\{ \left\| \sum_{n=0}^{\infty} b_n F(n)z^n \right\|_{L^p(\tilde{\Gamma}_1)} : F \in \mathcal{A}(\tilde{\Gamma}_1) \cap \mathcal{A}(\tilde{\Gamma}_2) \right\}.$$

If  $\tilde{\Gamma}_3$  is a third such curve, and  $b_1 \in M_p(\tilde{\Gamma}_1, \tilde{\Gamma}_2)$ ,  $b_2 \in M_p(\tilde{\Gamma}_2, \tilde{\Gamma}_3)$ , and  
 $b_3 \in M_p(\tilde{\Gamma}_1, \tilde{\Gamma}_3)$ , and

$$\|b_3\|_{M_p(\tilde{\Gamma}_1, \tilde{\Gamma}_3)} \leq \|b_2\|_{M_p(\tilde{\Gamma}_2, \tilde{\Gamma}_3)} \|b_1\|_{M_p(\tilde{\Gamma}_1, \tilde{\Gamma}_2)}$$

THEOREM 8. Let  $b \in l^\infty$  and  $f_b(n) = b(n) \exp(in\theta)$ . If  $f_b \in M_p(\tilde{\Gamma})$  for  
 some  $\tilde{\Gamma} \in \mathcal{M}$ , where  $\mathcal{M}$  is the set of all such curves, and if  $\alpha < \infty$ , then  
 $f_b \in M_p(\tilde{\Gamma})$ , and

$$\|f_b\|_{M_p(\tilde{\Gamma})} \leq \|b\|_{l^\infty} \|f_b\|_{M_p(\tilde{\Gamma})}.$$

The following example shows that we cannot expect the relation

$$\|f_b\|_{M_p(\tilde{\Gamma})} \leq \|b\|_{l^\infty} \|f_b\|_{M_p(\tilde{\Gamma})}$$
 to hold in general ([McQ3]). Although it does hold when the curve  $\tilde{\Gamma}$  is flat

with  $A = \tilde{\Gamma} = \{A(n) : A(0) > 0 \text{ and } n = \min A(n) < 0\}$ . For any integer  $S$   
 let  $b_S$  be the sequence in  $l^\infty$  defined by  $b_S(n) = 1$  if  $|n| \leq S$  and  $b_S(n) = 0$

References

[CM2] Au-delà des opérateurs pseudo-  
 [CMCM] S. Coifman, A. McIntosh et A.  
 opérateur borné sur  $L^p$  pour les cou-  
 (1982), 361-387.  
 [Dav1] G. David - Opérateurs intégraux sing-  
 liers, Ann. Sc. Ecole Norm. Sup. 17  
 [DJS] G. David, S. L. Journé et S. Semmes -  
 fonctions para-accrétes et interpolat-  
 ion, 1-56.  
 [FJR] E. B. Fabes, M. Jodeit, J. R. Ken-  
 nedy and J. Torrea - Boundary value problems on Sierpinski triangles,  
 [GQW] G. Gaudry, T. Qian and S.-L. Wan-  
 g - Singular integrals with holomorphic kernels on star-  
 shaped Lipschitz curves, Proc. Amer. Math. Soc. 121 (1994), 133-150.

[McQ] O. Lozano, A. McIntosh and T. Qian -  
 singular convolution operators on Lipschitz  
 surfaces, Illinois Math. Surv. 5 (1992), 45-48.  
 [L] L. L. Lorentz - On the boundedness of singular  
 integrals, Illinois Math. Surv. 1 (1956), 1-16.

[McQ3] Singular integrals along Lipschitz curves with holomorphic kernels.  
 Proc. Amer. Math. Soc. 121 (1994), 167-176.

- [Q4] T. Qian, *A holomorphic extension result*, Complex Variables Theory Appl. 32 (1996), 59–77.
- [Q5] —, *Singular integrals on star-shaped Lipschitz surfaces in the quaternionic space and generalizations to  $\mathbb{R}^n$* , in: Proc. Conf. on Clifford and Quaternionic Analysis and Numerical Methods, June 1996, to appear.
- [S] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, Princeton, N.J., 1970.
- [SW] E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, Princeton, N.J., 1971.
- [V] G. Verchota, *Layer potentials and regularity for the Dirichlet problems for Laplace's equation in Lipschitz domains*, J. Funct. Anal. 59 (1984), 572–611.
- [Z] A. Zygmund, *Trigonometric Series*, 2nd ed., Cambridge Univ. Press, London and New York, 1968.

Department of Mathematics  
The New England University  
Armidale, New South Wales 2351  
Australia  
E-mail: tao@neumann.une.edu.au

Received May 26, 1994

Revised version November 5, 1994

(3278)

called the maximum principle, says that a function holomorphic and of exponential type on a sector  $S$  of opening less than  $\pi$  is bounded if it is bounded on the boundary of  $S$ . The second one, ([L]), called the quasi-analyticity principle, says that a holomorphic function  $F$  on a sector  $S$  vanishes if the opening of  $S$  is greater than  $\pi$  and  $F$  is exponentially decreasing in  $S$ .

In the present paper we study the quasi-analyticity principle in the critical case of a half plane  $H$ . To ensure vanishing of  $F$  in that case we assume that  $F$  is of exponential type in  $H$  and decreases exponentially along the boundary of  $H$ . More precisely, we have

**THEOREM 1** (Quasi-analyticity principle, continuous version): Let  $F \in \mathcal{O}(\{\operatorname{Re} z > 0\}) \cap C^0(\{\operatorname{Re} z \geq 0\})$  be of exponential type, i.e.

$$(1) \quad |F(z)| \leq C e^{c|z|} \quad \text{for } \operatorname{Re} z \geq 0 \text{ with some } C < \infty \text{ and } c < \infty.$$

If

$$(2) \quad |F(\pm ir)| \leq C e^{c^\pm r} \quad \text{for } r \geq 0$$

with some  $c^\pm \in \mathbb{R}$  such that  $c^+ + c^- < 0$  then  $F \equiv 0$ .

The elementary proof of Theorem 1 is based on the Laplace integral representation of holomorphic functions of exponential type.

1991 Mathematics Subject Classification: 30D15, 44A15, 46F12, 46F20.

Key words and phrases: quasi-analyticity, Laplace distributions, Laplace ultradistri-

## A Phragmén–Lindelöf type quasi-analyticity principle

by

GRZEGORZ ŁYSIK (Warszawa)

**Abstract.** Quasi-analyticity theorems of Phragmén–Lindelöf type for holomorphic functions of exponential type on a half plane are stated and proved. Spaces of Laplace distributions (ultradistributions) on  $\mathbb{R}$  are studied and their boundary value representation is given. A generalization of the Painlevé theorem is proved.

**1. Introduction and statement of the main results.** The well-known Phragmén–Lindelöf theorem consists of two parts. The first one ([H]),